

# A simple inverter for polarization transformations

**Rajendra Bhandari**

March 3, 2010

Raman Research Institute,  
Bangalore 560 080, India.  
email: bhandari@rri.res.in

## Abstract

It is shown that a transparent birefringent element whose eigenstates are a pair of elliptical polarizations with principal axes along the  $x$  and  $y$  axes, sandwiched between a pair of orthogonal halfwave plates with principal axes along directions at 45 deg to the  $x$  and  $y$  axes, is equivalent to an element with the same eigenstates and eigenvalues but with the fast and the slow eigenstates interchanged. The device thus produces the inverse of the original unitary transformation. A similar result holds for a pure dichroic element. With electrically switchable halfwave plates such a device can be used to switch the sign of optical activity or to rotate through 90 deg, without any moving parts, a linear retarder or a linear polarizer.

OCIS Codes 260.5430, 230.4110, 230.2090, 280.4788

In this paper we point out an interesting application in polarization optics of some well known results in the theory of  $2 \times 2$  complex matrices. It is well known that any non-depolarizing polarization transformation can be represented by a  $2 \times 2$  complex matrix (Jones matrix) and that any such matrix  $S$  can be expressed as a product  $S = UD$  or  $S = D'U$ , where  $U$  is a unitary matrix and  $D, D'$  are nonunitary matrices with orthogonal eigenstates and real eigenvalues [1, 2]. We shall choose the first of the two decompositions and  $D$  will be called a “pure dichroic element”. The matrix  $U$  can be written as a product of an isotropic phase factor  $\exp(i\alpha)$  and a matrix with determinant  $+1$  whose eigenvalues are  $\exp(-i\delta)$  and  $\exp(i\delta)$ . Similarly the matrix  $D$  can be written as a product of an isotropic attenuation factor  $\exp(-\eta)$  and a matrix with unit determinant with eigenvalues  $\exp(-\gamma)$  and  $\exp(\gamma)$ . From now on we shall ignore the isotropic factors and assume that  $U$  and  $D$  are matrices with unit determinant, i.e. elements of the group  $SL(2, \mathbb{C})$ ; the former also being an element of the  $SU(2)$  subgroup of  $SL(2, \mathbb{C})$ . We note that for an arbitrary element of  $SL(2, \mathbb{C})$  the sets of orthogonal eigenstates of  $U$  and  $D$  are different.

Let us first consider a unitary transformation  $U$ . Let  $|u\rangle$  and  $|\tilde{u}\rangle$  be the two orthogonal eigenvectors of  $U$  with eigenvalues  $\exp(-i\delta)$  and  $\exp(i\delta)$  respectively. Let  $A$  be some other element of  $SU(2)$  such that  $A|u\rangle = |u'\rangle$  and  $A|\tilde{u}\rangle = |\tilde{u}'\rangle$ . It is then easy to verify that the matrix  $U' = AUA^\dagger$  has the states  $|u'\rangle$  and  $|\tilde{u}'\rangle$  as its eigenvectors, with eigenvalues  $\exp(-i\delta)$  and  $\exp(i\delta)$  respectively. The matrix  $A^\dagger$  is the hermitian conjugate of  $A$ , which is the same as  $A^{-1}$  when  $A$  is unitary. Now if the matrix  $A$  is chosen such that  $|u'\rangle = |\tilde{u}\rangle$  and  $|\tilde{u}'\rangle = |u\rangle$  then we have the result that the matrix  $AUA^\dagger$  has the states  $|\tilde{u}\rangle$  and  $|u\rangle$  as its eigenstates with eigenvalues  $\exp(-i\delta)$  and  $\exp(i\delta)$  respectively.  $U'$  therefore represents the transformation inverse of that represented by  $U$ . For example if  $U$  corresponds to pure optical activity then the matrix  $U' = AUA^\dagger$  also corresponds to pure optical activity of the same magnitude but with the sign changed. If on the other hand  $U$  represents a linear retarder with its fast axis oriented at an angle  $\psi$ , the matrix  $AUA^\dagger$  represents a linear retarder with the same retardation but with its fast axis oriented at  $\psi + 90^\circ$ .

It is not difficult to see what the matrix  $A$  is. A state  $|u\rangle$  represented by the point  $(\theta, \phi)$  on the Poincaré sphere can be taken to its orthogonal state  $|\tilde{u}\rangle$  by means of a  $\pi$  rotation about the point  $(90^\circ, \phi + 90^\circ)$ . This corresponds to a halfwave plate whose fast axis makes, in real space, an angle  $45^\circ$  with the major axis of the polarization ellipse corresponding to the state  $(\theta, \phi)$  on the

Poincaré sphere. This immediately leads to the first main result of the paper, i.e. a unitary polarization element  $U$  corresponding to a rotation through  $2\delta$  about a point  $(\theta, \phi)$  on the Poincaré sphere, sandwiched between two orthogonal halfwave plates whose fast axes make, in real space, angles  $\pm 45^\circ$  with the major axis of one of the ellipses representing the eigenpolarizations of  $U$  will correspond to a rotation through an angle  $-2\delta$  about the point  $(\theta, \phi)$  on the sphere. By orthogonal halfwave plates we mean those whose fast axes make an angle  $90^\circ$  with each other.

Let us next consider a dichroic transformation represented by a pure dichroic element  $D$ , i.e. an element with orthogonal polarization eigenstates  $|d\rangle$  and  $|\tilde{d}\rangle$  and eigenvalues  $\exp(-\gamma)$  and  $\exp(\gamma)$  respectively. It can be shown exactly as above that such an element, sandwiched between two orthogonal halfwave plates whose fast axes make angles  $\pm 45^\circ$  with the major axis of one of the ellipses representing the eigenpolarizations of  $D$  will correspond to a transformation which is inverse of  $D$ , i.e. have  $|\tilde{d}\rangle$  and  $|d\rangle$  as eigenstates with eigenvalues  $\exp(-\gamma)$  and  $\exp(\gamma)$  respectively;  $\gamma$  being a real quantity. If  $D$  corresponds to a linear polarizer, the sandwich corresponds to the same linear polarizer, rotated through  $90^\circ$ .

With the use of electro-optic modulators, i.e elements which can be switched electrically from being unit elements to being halfwave plates, the above considerations lead to the possibility of remotely inverting a unitary or a pure dichroic polarization transformation, for example (i) switching the sign of optical activity of an optically active sample, (ii) rotating the principal axes of a linear retarder through  $90^\circ$  or (iii) rotating through  $90^\circ$  the transmission axis of a linear polarizer, all the operations being without the use of any moving parts.

It is easy to show that the above device inverts not only unitary and pure dichroic transformations but also inverts a product  $S = UD$  of the two kinds of transformation if  $S$  and  $D$  have the same orthogonal pair of eigenstates. For then, since  $U$  and  $D$  commute,

$$\begin{aligned} S' &= ASA^\dagger = (AUA^\dagger)(ADA^\dagger) = U^{-1}D^{-1} \\ &= (DU)^{-1} = (UD)^{-1} = (S)^{-1} \end{aligned} \tag{1}$$

The above equation implies that if a linear polarizer has a small amount of linear birefringence with the same principal axes as those of the polarizer, the device described above would still work, i.e. the imperfect polarizer,

sandwiched between a pair of orthogonal halfwave plates at  $\pm 45^\circ$  is equivalent to the same imperfect polarizer, rotated through  $90^\circ$ .

While the result described in this paper is trivial as a mathematical result we have been unable to find in literature an instance of its use in polarization optics, hence this communication.

## References

- [1] Wu-Ki Tung, *Group theory in physics*, (World Scientific 1985), Ch.10.
- [2] S. Y. Lu and R. A. Chipman, “Homogenous and inhomogenous Jones matrices”, J. Opt. Soc. Am. A **11**, 766 (1994).